



## Research Article

# On Characterization of Various Finite Subgroups of Abelian Groups

N. B. Okelo\*

School of Mathematics and Actuarial Science,  
Jaramogi Oginga Odinga University of Science and Technology,  
P. O. Box 210-40601, Bondo-Kenya.

\*Corresponding author's e-mail: [bnyaare@yahoo.com](mailto:bnyaare@yahoo.com)

## Abstract

The present paper, we characterize finite subgroups. Throughout  $G$  always denote a finite group. Let  $H$  be a subgroup group of  $G$ . We have  $H \geq H \cap H^x \geq 1$ , for any  $x \in G$ . We call  $H$  to be a TI-subgroup of  $G$  if  $H \cap H^x = H$  or  $1$  for any  $x \in G$ . We have shown that if  $H$  is normal in  $G$  or if  $H$  is of a prime order, then  $H$  is a TI-subgroup.

**Keywords:** Group; Finite group; TI-subgroup; Abelian group.

## Introduction

Characterizations in group theory regarding subgroups have been done over a period of time by many authors [1-5]. A topic of some interest is to investigate the finite groups in which certain subgroups are assumed to be TI-subgroups. The author in [6] classified the finite groups all of whose subgroups are TI-subgroups. In [7, 8], Guo, they classified the finite groups whose abelian subgroups are TI-subgroups. The aim of this paper is to study the finite AQTI-groups, that is, all of whose abelian subgroups are QTI (that means quasi-trivialintersection)-subgroups [9]. We obtain a classification of the AQTI-groups in Theorem 3.3 (nilpotent case) and Theorem 3.7 (non-nilpotent case). The aim of this work is to characterize TI, ATI and QTI subgroups in depth.

## Research Methodology

### Definition 1.1

A subgroup  $H$  of  $G$  is called a QTI-subgroup if  $C_G(x) \leq N_G(H)$  for any  $1 \neq x \in H$ .

Clearly a TI-subgroup is a QTI-subgroup. However, the converse is not true [10].

### Example 1.2

Let  $V$  be an elementary abelian 3-group of order 35 and  $H$  be a subgroup of  $GL(5, 3)$  of order

$11^2$ . Let  $G = HV$ , where  $H$  acts on  $V$  in a natural way. Since 11 does not divides  $3^a - 1$  for any  $a < 5$ , the actions of  $H$  and its nonidentity subgroups on  $V$  are irreducible and fixed-point-free. It follows that  $N_G(W) = V$  for any proper subgroup  $W$  of  $V$  and that  $C_G(w) = V$  for any  $1 \neq w \in W$ , and therefore  $W$  is a QTI-subgroup of  $G$ . In fact, it is not difficult to see that all abelian subgroups of  $G$  are QTI-subgroups, and therefore  $G$  is an AQTI-group. Let  $W_0$  be a subgroup of  $V$  of order  $3^4$ . Since  $|W_0 \cap W_0^x| = 3^3$  for any  $1 \neq x \in H$ ,  $W_0$  is not a TI-subgroup. A very important question to ask at this juncture is: Under which additional condition  $P$ , a QTI-subgroup is necessary a TI-subgroup? that is, QTI-subgroup+  $P \Rightarrow$  TI-subgroup.

## Results and discussion

### Lemma 3.1

Let  $G$  be an AQTI-group. Then the following statements hold.

- (i) Any subgroup of  $G$  is again an AQTI-group.
- (ii) For any abelian subgroup  $H$  of  $G$ , if  $H \cap Z(G) > 1$ , then  $H$  is normal in  $G$ .
- (iii) For any  $1 \neq x \in G$ ,  $C_G(x)$  is nilpotent.

Proof : (i) and (ii) are clear. (iii) For any cyclic subgroup  $A/\langle x \rangle$  of  $C_G(x)/\langle x \rangle$ , A is an abelian subgroup of an AQTG group  $C_G(x)$ , and so A is normal in  $C_G(x)$  (see (2)). It follows that all cyclic subgroups (and so all subgroups) of  $C_G(x)/\langle x \rangle$  are normal in  $C_G(x)/\langle x \rangle$ . Then  $C_G(x)/\langle x \rangle$  is nilpotent, and so  $C_G(x)$  is nilpotent. Recall that a CN-group is a group in which the centralizer of any nonidentity element is nilpotent. Now the above lemma implies that an AQTG group is a CN-group. For any finite group G, we define its prime graph  $\Gamma(G)$  (see [8]) as follows: Whose vertex set is  $\pi(G)$ , and two vertices p, q are jointed by an edge if G has an element of order pq. If  $\sigma$  is a vertex set of a connected component of  $\Gamma(G)$ , then  $\sigma$  is called a prime component of G. This completes the proof.

### Lemma 3.2

([2, Theorem 2.2]) Let G be a CN-group and  $\sigma$  a prime component of G. Then G possesses a nilpotent Hall  $\sigma$ -subgroup H, and any  $\sigma$ -subgroup is contained in some G-conjugate of H, furthermore H is a TI-subgroup if in addition  $|\sigma| \geq 2$ . In particular, if G is a nonnilpotent AQTG-group, then  $\Gamma(G)$  is disconnected.

We note that the original proof of above lemma is elementary. Recall that a Hamiltonian group is a nonabelian group in which all subgroups are normal. It is known that a Hamiltonian group is a direct product of  $Q_8$ , an elementary abelian 2-group and an abelian group of odd order. For a p-group G, we put  $V_1(G) = \langle x^p \mid x \in G \rangle$ .

### Theorem 3.3

For a finite p-group G, the following statements are equivalent.

- (1) All subgroups of G are TI-subgroups.
- (2) All abelian subgroups of G are TI-subgroups.
- (3) All abelian subgroups of G are QTI-subgroups, ie., G is an AQTG-group.
- (4) G is one of the following p-groups:
  - (4.1) G is an abelian p-group.
  - (4.2) G is a Hamiltonian 2-group, that is a product of  $Q_8$  and an elementary abelian 2-group.

(4.3) G is the central product of  $Q_8$  and  $D_8$ ;

(4.4)  $G/Z(G)$  is of order  $p^2$ ,  $Z(G)$  is cyclic and  $G' \cong Z_p$  is the only minimal normal subgroup of G.

Remark: The objective of the paper [6] is to show the following: The finite p-groups all of whose abelian subgroups are TI-subgroups, are just the groups of types (4.1)-(4.4). Our arguments (of Theorem 3.3) are much shorter than those in [6].

Proof : We need only to show (3) implying (4). Suppose that all abelian subgroups of G are normal. Then all subgroups of G are normal, and so G is of type (4.1) or type (4.2). In what follows we assume that G has an abelian but not normal subgroup, and we will show that G is of type (4.3) or type (4.4). Observe first that for any nontrivial abelian subgroup A of G, A is normal in G iff  $A \cap Z(G) > 1$  (see Lemma 3.1(ii)).

Step 1.  $Z(G)$  is cyclic. Suppose that  $Z(G)$  is not cyclic and let A be any abelian subgroup of G. If  $A \cap Z(G) > 1$ , then A is normal in G. If  $A \cap Z(G) = 1$ , then  $AU, AV$  are normal in G where  $U, V \cong Z_p$  are distinct subgroups of  $Z(G)$ , and so  $A = AU \cap AV$  is normal. This implies that all abelian subgroups are normal, which contradicts our assumption.

Step 2. Let Z be the unique minimal normal subgroup of G. Then  $G/Z$  is abelian, and  $Z = G'$ . Let  $A/Z$  be any cyclic subgroup of  $G/Z$ . Then A is normal in G because A is abelian with  $A \cap Z(G) \geq Z$ . It follows that all subgroups of  $G/Z$  are normal. Suppose  $G/Z$  is nonabelian. Then G is a Hamiltonian 2-group, and so  $G/Z \cong Q_8 \times Z_2 \times \dots \times Z_2$ . Let  $T/Z \cong Q_8$ . Clearly T is normal in G and so  $T'$  is normal in G. Since Z is the unique minimal normal subgroup of G,  $T' \geq Z$ , and this implies that  $|T/T'| = 4$ . Now applying [3, Ch3, theorem, 11.9], we conclude that  $Z(T) = Z$ . By [3, Page 94, exercise 58], we get a contradiction. Thus  $G/Z$  is abelian, and so  $Z = G'$ .

Step 3. Final part of proof. Since  $G' = Z$  is the unique minimal normal subgroup of G, it follows by [5, Lemma 12.3] that  $G/Z(G)$  is elementary abelian and that all nonlinear irreducible

complex characters of  $G$  have degree  $\sqrt{|G/Z(G)|}$ .

Since  $G$  has an abelian but not normal subgroup  $A$  and  $A \cap Z(G) = 1$ , we can find an element  $t$  such that  $\langle t \rangle \cap Z(G) = 1$ . Then  $H =: C_G(t) < G$ .

..... It is easy to see that  $H$  is a maximal subgroup of  $G$  and that all abelian subgroups of  $H$  are normal (and so  $H$  is abelian or  $H = Q_8 \times Z_2 \times \dots \times Z_2$ ). Suppose that  $H$  is abelian. Since  $|G : H| = p$ , all nonlinear irreducible complex characters of  $G$  have degree  $p$ , and this implies that  $|G/Z(G)| = p^2$ , thus  $G$  is of type (4.4). Suppose that  $H = Q_8 \times Z_2 \times \dots \times Z_2$ . Then  $G$  possesses an abelian subgroup of index 4. It follows that all nonlinear irreducible complex characters of  $G$  have degree 2 or 4. Thus either  $|G/Z(G)| = 4$  and then  $G$  is of type (4.4), or  $|G/Z(G)| = 2^4$ . Let us investigate the case when  $|G/Z(G)| = 2^4$ . For this case, ..... we can prove that  $G$  is an extra special 2-group of order  $2^5$  (Thus,  $G \cong D_8 * D_8$  or  $D_8 * D_8$ ) and that the case  $G = D_8 * D_8$  is impossible. And hence  $G$  is a central product of  $D_8$  and  $Q_8$ , ie.,  $G$  is of type (4.3).

### Lemma 3.4

Let  $G$  be a finite group. Then  $G$  is an AQTI-subgroup iff  $G$  satisfies the following conditions:

(1)  $G$  is a CN-group,

(2) Let  $\sigma$  be any prime component of  $G$  and let  $M$  be a Hall  $\sigma$ -subgroup of  $G$ . Then either  $M$  is one of the  $p$ -groups listed in theorem 3.3, or  $M$  is abelian, or  $M$  is a Hamiltonian group.

Applying Theorem 3.3 and Lemma 3.4, we obtain the following result.

### Theorem 3.5

Let  $G$  be a nilpotent group. Then  $G$  is an AQTI-group if and only if one of the following holds.

(1)  $G$  is abelian.

(2)  $G$  is a Hamiltonian group.

(3)  $G$  is of type (4.3) or (4.4) in Theorem 3.1.

The proof of Lemma 3.4: Suppose that  $G$  is an AQTI-group. By Lemma 2.2,  $G$  is a CN-group, and  $G$  possesses a nilpotent Hall  $\sigma$ -subgroup  $M$  for any prime component  $\sigma$  of  $G$ . Clearly  $M$  is again an AQTI-subgroup, and we need to show that if  $|\sigma| \geq 2$  then all subgroups of  $M$  are

normal in  $M$ . Assume this is not true. Write  $M = P \times Q$ , where  $Q$  is a nontrivial  $p$ -group, and  $P \in Syl_p(M)$  has an abelian but not normal subgroup  $P_1$ . Let  $1 \neq x \in Z(Q) \leq P_1 \times Q$ . As  $P_1 \times Z(Q)$  is a QTI-subgroup of  $M$ ,  $M = C_M(x) \leq N_M(P_1 \times Z(Q)) = N_P(P_1) \times Q$ , and this implies that  $P_1$  is normal in  $P$ , a contradiction. Suppose conversely that  $G$  satisfies the conditions of Lemma 3.2. Let  $H$  be an abelian subgroup of  $G$  and  $1 \neq x \in H$ . Let  $p$  be a prime divisor of  $|H|$  and let  $\sigma$  be a prime component containing  $p$  of  $G$ . By Lemma 2.2 we may assume  $C_G(x) \leq M$ . If  $|\sigma| \geq 2$ , then  $M$  is a Hamiltonian group or an abelian group, thus  $H$  is normal in  $M$ , and so  $C_G(x) = C_M(x) \leq M = N_M(H) \leq N_G(H)$ . If  $|\sigma| = 1$ , then  $M$  is an AQTI-group of prime power order, so  $C_G(x) = C_M(x) \leq N_M(H) \leq N_G(H)$ . Thus  $H$  is a QTI-subgroup of  $G$ , and therefore  $G$  is an AQTI-group. If  $G = HN$  is a Frobenius group with a kernel  $N$  and a complement  $H$ , then we say that  $H$  acts frobeniusly on  $N$ . In this case, we know that  $N$  is nilpotent and any Sylow subgroup of  $H$  is either a cyclic group or a generalized quaternion group, and that  $\pi(H)$ ,  $\pi(N)$  are just two prime components of  $G$  (see [8]). If there are  $M, N < G$  such that  $G/N$  is a Frobenius group with  $M/N$  as its kernel and  $M$  is a Frobenius group with  $N$  as its kernel, then  $G$  is called a 2-Frobenius group, and such a 2-Frobenius group is denoted by  $\text{Frob}_2(G, M, N)$ . In this case, we know that  $G$  is solvable, and that  $\pi(M/N)$  and  $\pi(G/M) \cup \pi(N)$  are just two prime components of  $G$  (see [8]).

### Lemma 3.6

Let  $G = HN$  be a Frobenius group with a complement  $H$  and a kernel  $N$ . If  $G$  is an AQTI-group, then the following statements hold.

(1)  $H$  is either a cyclic group or a product of  $Q_8$  with a cyclic group of odd order.

(2)  $N$  is either an abelian group or of type (4.4) listed in Theorem 3.3.

Proof: Since  $G$  is a Frobenius group,  $\Gamma(G)$  has just two connected components with  $\pi(H)$ ,  $\pi(N)$  as its vertex sets.

(1) If  $H$  is nonnilpotent, then Lemma 2.2 implies that  $\Gamma(H)$  is disconnected, and then  $\Gamma(G)$  has at least three connected components, a contradiction. Thus  $H$  is nilpotent. If  $P \in Syl(H)$  is not cyclic, then  $P$  is a generalized quaternion group, and then  $P \cong Q_8$  by Theorem 3.1. The result follows.

(2) Since  $N$  is the Frobenius kernel,  $N$  is nilpotent. Assume that  $N$  is nonabelian and let  $P$  be a nonabelian Sylow  $p$ -subgroup of  $N$ . Then  $P$  is one of the three types listed in Theorem 3.1. Assume that  $P \cong Q_8 \times Z_2 \times \dots \times Z_2$ . Then  $V_1(P)$  is a normal subgroup of  $G$  of order 2, which is clearly impossible. Assume that  $P$  is the central product of  $Q_8$  and  $D_8$ . Then  $Z(P)$  lies in  $Z(G)$ , a contradiction. Thus  $P$  is of type (4.4) in Theorem 3.3, and then  $N = P$  by Theorem 3.3.

### Lemma 3.7

Let  $G = \text{Frob}_2(G, H, K)$ . If  $G$  is an AQTI-subgroup, then  $G$  is isomorphic to symmetric group  $S_4$ .

Proof: Note that  $G$  is solvable with just two prime components  $\pi_1 = \pi(H/K)$  and  $\pi_2 = \pi(G) - \pi_1$ , and that  $G$  has a nilpotent Hall  $\pi_2$ -subgroup  $W$  (see Lemma 3.2). Clearly  $K$  is the Fitting subgroup of  $G$ , thus  $C_w(K) \leq C_G(K) \leq K$ , and so  $W > K > Z(W)$ . Let  $p \in \pi(G/H)$  and  $P$  be a Sylow  $p$ -subgroup of  $W$ . Since  $K > Z(W) \geq Z(P)$ ,  $P \cap K \geq Z(P)$  is nontrivial. Let  $G_1 > P$  be a  $\pi_1 \cup \{p\}$ -Hall subgroup of  $G$ . It follows that  $G_1 = \text{Frob}_2(G_1, H \cap G_1, P \cap K)$ . Assume that  $G_1 < G$ . Then induction yields that  $G_1 \cong S_4$ , thus  $P \in \text{Syl}_2(S_4)$  is isomorphic to  $D_8$ , and then  $W = P$  by Theorem 3.3, so  $G_1 \cong S_4$  as wanted. In what follows, we assume that  $\pi_2 = \{p\}$ . Then  $W$  is one of the nonabelian  $p$  groups listed in Theorem 3.3.

Case 1. Assume that  $W \cong Q_8 \times Z_2 \times \dots \times Z_2$ . As  $W > K > Z(W)$ ,  $K$  is a product of  $Z_4$  and an elementary abelian 2-group. It follows that  $V_1(K) < G$  with  $|V_1(K)| = 2$ , a contradiction.

Case 2. Assume that  $W$  is the central product of  $Q_8$  and  $D_8$ . As  $W > K > Z(W)$ ,  $|K| \in \{4, 8, 16\}$ . If  $K$  is abelian, then  $K \in \{Z_4 \times Z_2, Z_4, Z_2 \times Z_2\}$  (see [3, Ch3, Theorem 13.8]). Now  $K/\Phi(K) = Z_2$  or  $Z_2 \times Z_2$ , it follows that  $G/K \leq \text{Aut}(K/\Phi(K)) \leq S_3$ , then  $|P| \leq 16$ , a contradiction. If  $K$  is nonabelian and of order 16, then  $K \cong Q_8 \times Z_2$  or  $|K/Z(K)| = 4$  with  $Z(K) \cong Z_4$ . For the first case, let  $Z = V_1(K)$ ; and for the second case, let  $Z = V_1(Z(K))$ . Then  $Z$  is normal in  $G$  with  $|Z| = 2$ , a contradiction. If  $K$  is nonabelian and of order 8, then  $K \cong Q_8$  or  $D_8$ , and then  $G/K \leq \text{Aut}(K/\Phi(K)) = \text{Aut}(Z_2 \times Z_2) = S_3$ , thus  $|P| = 16$ , a contradiction.

Case 3. Assume that  $W/Z(W) \cong Z_p \times Z_p$  and  $Z(W)$  is cyclic. Then  $K$  is abelian with  $|W : K| = |K : Z(W)| = p$ . Note that  $G = N_G(U)H = N_G(U)K$  by Frattini argument, where  $U$  is a Hall  $\pi_1$ -subgroup of  $G$ . Clearly  $N_G(U) \cap K = N_K(U) = 1$ , and so  $N_G(U) \cong G/K$  is a Frobenius group with a complement of order  $p$ . Suppose  $K$  is not elementary abelian. Then  $V_1(K)$  is a nontrivial cyclic normal subgroup of  $G$ . Let us consider  $G_1 = N_G(U)V_1(K)$ . We see that  $V_1(K) = \text{Fit}(G_1)$ , and  $N_G(U) \leq \text{Aut}(V_1(K))$  is abelian, a contradiction. Hence  $K$  is elementary abelian, and in particular  $Z(W) \cong Z_p$ . Now  $N_G(U) \leq \text{Aut}(K) = \text{Aut}(Z_p \times Z_p) = \text{GL}(2, p)$ . Note that if  $p > 2$ , then it is easy to check that  $\text{GL}(2, p)$  has no subgroup which is a Frobenius group with a complement of order  $p$ . This implies that  $K \cong Z_2 \times Z_2$ , and hence  $N_G(U) \cong S_3$ , and  $G \cong S_4$ .

### Theorem 3.8

Let  $G$  be a nonnilpotent group. Then  $G$  is an AQTI-subgroup iff  $G$  is one of the following groups.

(1)  $G = HN$  is a Frobenius group with a complement  $H$  and a kernel  $N$ , where  $N$  is abelian, and  $H$  is either a cyclic group or a product of  $Q_8$  with a cyclic group of odd order.

(2)  $G = HN$  is a Frobenius group with a complement  $H$  and a kernel  $N$ , where  $H$  is a cyclic subgroup of  $Z_{p-1}$  and  $N$  is a  $p$ -group of the type (4.3) in Theorem 3.3.

(3)  $G \cong S_4$ .

(4)  $G \cong L_2(q)$ , where  $q = 5, 7, 9$ .

Proof: Suppose that  $G \in \{S_4, L_2(5), L_2(7), L_2(9)\}$ . Then it is easy to check that  $G$  is an AQTI-group. Suppose that  $G$  is a Frobenius group of type (1) or (2). We also conclude by Lemma 3.2 that  $G$  is an AQTI-group. Suppose that  $G$  is a nonnilpotent AQTI-group. Then the prime graph  $\Gamma(G)$  is disconnected (see Lemma 3.2). Assume that  $G$  is solvable. It is well known that  $G$  is a Frobenius or 2-Frobenius group (see [8]), and then Lemma 3.6 and Lemma 3.7 imply that  $G$  is of type (1) or type (2). In what follows, we assume that  $G$  is a nonsolvable AQTI-group. Let  $N = \text{Sol}(G)$ , the maximal normal solvable subgroup of  $G$ . It follows by [8] that  $G$  has a normal series  $N < H < G$  such that  $N$  and  $G/H$  are  $\pi$ -groups and  $H/N$  is a nonabelian simple group, where  $\pi$  is the prime component of  $G$  containing 2. Furthermore,  $N = \text{Sol}(G) = \text{Fit}(G)$ ,  $G/N \leq \text{Aut}(H/N)$ . Let  $P_1$  be a nilpotent Hall  $\pi$ -subgroup of  $G$  (see Lemma 3.2), and  $P = P_1 \cap H$ .

Claim 1. If  $N > 1$ , then  $\pi = \{2\}$ . Suppose that  $N > 1$  and  $|\pi| \geq 2$ . By Lemma 3.2,  $P_1$  is a TI-subgroup of  $G$ . Since  $N \leq P_1$  is a nontrivial normal subgroup of  $G$ ,  $P_1$  is normal in  $G$ , so  $G$  is solvable, a contradiction. Thus  $|\pi| = 1$  and so  $\pi = \{2\}$ .

Claim 2.  $N = 1$ . Suppose that  $N > 1$  and let  $E$  be any normal subgroup of  $G$  with  $1 < E \leq N$ . By claim 1,  $\pi = \{2\}$  and  $P$  is a 2-group. Assume that  $C_G(E)N > N$ . Since  $H/N$  is simple and is a unique minimal normal subgroup of  $G/N$ ,  $C_G(E)N \geq H$ . Then any odd order subgroup of  $H$  acts trivially on  $E$ , which is clearly impossible. Hence  $C_G(E) \leq N$ , and in particular  $P > N > Z(P)$ . Now  $P$  is one of the 2-groups listed in Theorem 3.1. Arguing as in the proof of Lemma 4.2, we can find a normal subgroup  $E$  of  $G$  with  $1 < E \leq N$  and  $E \leq Z_2 \times Z_2$ . It follows that  $G/C_G(E) \leq \text{Aut}(E)$  is solvable, and so  $G/N$  is solvable because  $C_G(E) \leq N$ , a contradiction.

Claim 3.  $H \cong L_2(q)$ , where  $q = 5, 7, 9$ . As  $N = 1$ ,  $H$  is a nonabelian simple group. Since  $H$  is an AQTI-group, by Lemma 3.1(iii)  $H$  is a CN-group. Note that the only simple nonabelian CN-groups are  $Sz(q)$ ,  $L_3(4)$ ,  $L_2(9)$ , and  $L_2(p)$  where  $p$  is a Fermat or a Mersenne prime (see [4, ChXI, Remark 3.12]). Assume that  $H \cong Sz(q)$ . Then  $|P| = q^2$ ,  $q = 22m+1$ , where  $P_0 = Z(P) = \{1\}$  is an elementary abelian group of order  $q$ . Checking the 2-groups listed in Theorem 3.1, we get a contradiction. Assume that  $H \cong L_3(4)$ . Then  $|P| = 2^6$ , and  $Z(P) \cong Z_2 \times Z_2$ . Checking the 2-groups listed in Theorem 3.3, we get a contradiction. Assume that  $H \cong L_2(p)$ , where  $p$  is a prime and  $p = 2^m + 1$  or  $2^m - 1$ . Then  $P$  is a dihedral group of order  $2^m$  (see [3, ChII, Theorem 8.27]). Checking the 2-groups listed in Theorem 3.3, we conclude that  $P \cong Z_2 \times Z_2$  or  $D_8$ . Thus either  $p = |P| + 1 = 5$  and then  $H \cong L_2(5)$ , or  $p = |P| - 1 = 7$  and then  $H \cong L_2(7)$ .

Claim 4.  $G = H \cong L_2(q)$ , where  $q = 5, 7, 9$ . It suffices to show that  $G = H$ . Otherwise,  $H < G \leq \text{Aut}(H)$ . We will apply [1] to get a contradiction. Assume that  $H \cong A_5$  (or  $L_2(7)$ ). Then  $G \cong S_5$  (or  $\text{PGL}(2, 7)$ ) has an element of order 6, so 2, 3 lie in the same prime component of  $G$ . However neither  $S_5$  nor  $\text{PGL}(2, 7)$  has a nilpotent Hall  $\{2, 3\}$ -subgroup, a contradiction. Assume that  $H \cong L_2(9)$ . Then  $G$  contains a subgroup which is isomorphic to  $L_2(9) : 2_1$ ,  $L_2(9) : 2_2$  or  $L_2(9) : 2_3$  (see [1]). If  $L_2(9) : 2_1 \leq G$ , then  $G$  has an element of order 6

but has no nilpotent Hall  $\{2, 3\}$ -subgroup, a contradiction. If  $L_2(9) : 2_2 \leq G$ , then  $G$  has an element of order 10 but has no nilpotent Hall  $\{2, 5\}$ -subgroup, a contradiction. If  $L_2(9) : 2_3 \leq G$ , then a Sylow 2-subgroup  $U$  of  $L_2(9) : 2_3$  has order 16 and  $|Z(U)| = 2$ , and we also get a contradiction by checking the 2-groups listed in Theorem 3.3. Thus  $G = H$  as desired.

## Conclusions

We conclude this paper by asking two important questions: Let  $H$  be a subgroup of a finite group  $G$ . Clearly,  $H \geq H \cap H^x \geq H_G = \cap_{x \in G} H^x \geq 1$ . We call  $H$  is a CTI-subgroup of  $G$  if  $H \cap H^x = H$  or  $H_G$  for any  $x \in G$ . Our question is to classify the finite  $p$ -groups (or finite groups) all of whose subgroups (or abelian subgroups) are CTI-subgroups. Secondly, what can we say about the finite groups with no nontrivial TI-subgroup. Here a trivial TI-subgroup is a normal subgroup or a subgroup of prime order.

## Conflicts of interest

Authors declare no conflict of interest.

## References

- [1] Conway JH, Curtis RT, Norton SP, Parker RA, Wilson RA. *Atlas of finite groups*, Oxford University Press (Clarendon), Oxford and New York; 1985.
- [2] Guo X, Li S, Flavell P. Finite groups whose abelian subgroups are TI-subgroup. *J Algebra* 2007;307:565-69.
- [3] Huppert B, Endlich GI. *Springer-Verlag*, Berlian Heidelberg, New York; 1979.
- [4] Huppert B, Blackburn N. *Finite groups III*, Springer-Verlag, Berlin Heidelberg, New York; 1982.
- [5] Isaacs IM. *Character Theory of Finite Groups*, Academic Press, New York; 1976.
- [6] Li S, Guo X. Finite  $p$ -groups whose abelian subgroups have a trivial intersection. *Acta Math. Sin.* 2007;23(4):731-34.
- [7] Walls G. Trivial intersection groups. *Arch Math* 1979;32:1-4.
- [8] Williams JS. Prime graph components of finite groups *J Algebra* 1981;69:487-13.
- [9] Vijayabalaji S, Sathiyaseelan N. Interval-Valued Product Fuzzy Soft Matrices and

its Application in Decision Making, Int J Mod Sci Technol 2016;1(7):159-63.

[10] Chinnadurai V, Bharathivelan K. Cubic Ideals in Near Subtraction Semigroups. Int J Mod Sci Technol 2016;1(8):276-82.

\*\*\*\*\*