

Research Article

Characterization of Properties of Aluthge Transforms in Banach Algebras

I. O. Okwany, N. B. Okelo, Judith J. E. J Ogal

School of Mathematics, Statistics and Actuarial Science,
Jaramogi Oginga Odinga University of Science and Technology,
P. O. Box 210-40601, Bondo-Kenya.

*Corresponding author’s e-mail: bnyaare@yahoo.com

Abstract

Let \mathcal{H} be a complex separable Hilbert space with and let $\mathcal{B}(\mathcal{H})$ be the Banach algebra of all bounded linear operators on \mathcal{H} . In the present paper we characterize Aluthge transforms in Banach algebras. We considered classical and maximal numerical ranges of these transforms and finally we give their relationships.

Keywords: Hilbert space; Numerical ranges; Aluthge transforms; Banach algebra.

Introduction

It is known that the numerical range $W(T)$ of T is the subset $W(T) = \{\langle Tx, x \rangle : x \in \mathcal{H}, \|x\| = 1\}$ of the complex plane \mathbb{C} [1]. It is known that $W(T)$ is always convex and the closure $\overline{W(T)}$ of $W(T)$ contains $\sigma(T)$. On the other hand, essential numerical range of T is the subset $W_e(T) = \{\lambda \in \mathbb{C} : \text{there exists a unit vector sequence } \{x_n\} \subset \mathcal{H} \text{ such that } x_n \text{ converges weakly to } 0, \langle Tx_n, x_n \rangle \rightarrow \lambda\}$. It is known [2] that $W_e(T)$ is also always non-empty closed and convex and contains $\sigma_e(T)$. In [3], we also have $W_e(T) = \bigcap \{W(T + K) : K \in \mathcal{K}(\mathcal{H})\}$.

In [4] the author introduced the concept of the maximal numerical range $W_0(T)$ of T to consider the norm of a derivation on $\mathcal{B}(\mathcal{H})$. The maximal numerical range of T is defined to be the subset $W_0(T) = \{\lambda \in \mathbb{C} : \text{there exists a unit vector sequence } \{x_n\} \subset \mathcal{H} \text{ such that } \langle Tx_n, x_n \rangle \rightarrow \lambda, \|Tx_n\| \rightarrow \|T\|\}$. It was proved in [5] that $W_0(T)$ is a non-empty closed and convex subset of \mathbb{C} . We note that $W_0(T)$ does not have translation property by scalar, that is $W_0(T + \lambda) \neq W_0(T) + \lambda$. In particular, we know that for any $\lambda_1 \neq \lambda_2$ in \mathbb{C} , $W_0(T + \lambda_1) \cap W_0(T + \lambda_2) = \emptyset$. For a more detailed discussion of the maximal numerical range we refer to [6]. For a subset Δ of \mathbb{C} , we denote by Δ^\wedge the closed convex hull of Δ . If $T \in \mathcal{B}(\mathcal{H})$ with a polar decomposition $T = U|T|$,

then the Aluthge transform \tilde{T} and *-Aluthge transform $\tilde{T}^{(*)}$ are defined by $\tilde{T} = |T|^{1/2} U |T|^{1/2}$ and $\tilde{T}^{(*)} = |T^*|^{1/2} U |T^*|^{1/2}$ respectively [7]. Note that both \tilde{T} and $\tilde{T}^{(*)}$ are independent of the choice of the partial isometry U in the polar decomposition of T . Recently, T, \tilde{T} and $\tilde{T}^{(*)}$ have been studied by many authors [8]. In this note, we consider the essential numerical range and the maximal numerical range of T, \tilde{T} and $\tilde{T}^{(*)}$. We prove that $W_e(\tilde{T}) = W_e(\tilde{T}^{(*)}) \subseteq W_e(T)$ and $W_0(\tilde{T} + \lambda) = W_0(\tilde{T}^{(*)} + \lambda)$ for all $\lambda \in \mathbb{C}$.

Let $T \in \mathcal{B}(\mathcal{H})$ and $T = U|T|$ be the polar decomposition of T , then [9] we have $N(T) = N(|T|) = N(U)$. In terms of the orthogonal

decomposition $\mathcal{H} = N(T) \oplus N(T)^\perp$ of \mathcal{H} , T has

$$T = \begin{pmatrix} 0 & A \\ 0 & B \end{pmatrix}$$

the following matrix form for some bounded linear operators A from $N(T)^\perp$ to $N(T)$ and B on $N(T)^\perp$. Now it is known [10] that

$$U = \begin{pmatrix} 0 & U_1 \\ 0 & U_2 \end{pmatrix} \text{ and } T^*T = \begin{pmatrix} 0 & 0 \\ 0 & A^*A + B^*B \end{pmatrix}$$

for some operators U_1 and U_2 . By a simple calculus, \tilde{T} has the following matrix

$$\tilde{T} = \begin{pmatrix} 0 & 0 \\ 0 & X \end{pmatrix}, \text{ where } X = (A^*A + B^*B)^{1/4} U_2 (A^*A + B^*B)^{1/4} \text{ on } N(T)^\perp.$$

It is known that $\tilde{T}^{(*)} = U\tilde{T}U^*$ and $\tilde{T} = U^*\tilde{T}^{(*)}U$. Note that U is a unitary operator from $N(T)^\perp$ to $N(T^*)^\perp$, then there is a unitary operator U_0 from $N(T)^\perp$ onto $N(T^*)^\perp$ such that $U = \begin{pmatrix} 0 & 0 \\ 0 & U_0 \end{pmatrix}$ from $N(T) \oplus N(T)^\perp$ to $N(T^*) \oplus N(T^*)^\perp$. It follows that $\tilde{T}^{(*)} = \begin{pmatrix} 0 & 0 \\ 0 & Y \end{pmatrix}$ with respect to the space decomposition $\mathcal{H} = N(T^*) \oplus N(T^*)^\perp$, where $Y = U_0 X U_0^*$.

Research methodology

Proposition 2.1

Let $T \in \mathcal{B}(\mathcal{H})$. Then for all $K \in \mathcal{K}(\mathcal{H})$, $T + K - \tilde{T} \in \mathcal{K}(\mathcal{H})$.

Proof

Since $(T + K)^*(T + K) = T^*T + K_1$ for some $K_1 \in \mathcal{K}(\mathcal{H})$, we have $|T + K|^2 = |T|^2 + K_1$. It follows that $(\pi(|T + K|))^2 = (\pi(|T|))^2$ and $\pi(|T + K|) = \pi(|T|)$, which implies that $|T + K| - |T| \in \mathcal{K}(\mathcal{H})$ and again $|T + K|^{1/2} - |T|^{1/2} \in \mathcal{K}(\mathcal{H})$.

Let $T = U|T|$ and $T + K = V|T + K|$ be the decomposition of T and $T + K$ respectively. Note that $K = V|T + K| - U|T| \in \mathcal{K}(\mathcal{H})$, then $V(|T + K| - |T|) + (V - U)|T| \in \mathcal{K}(\mathcal{H})$, and therefore $(V - U)|T| \in \mathcal{K}(\mathcal{H})$ since $V(|T + K| - |T|) \in \mathcal{K}(\mathcal{H})$. If $|T|$ is invertible, then $(V - U)|T|^{1/2} \in \mathcal{K}(\mathcal{H})$. Otherwise, put $f(t) = t^{1/2}$, $t \in [0, |T|]$. We may choose a sequence of polynomials $P_n(t)$ with $P_n(0) = 0$ such that $\lim_{n \rightarrow \infty} \|P_n - f\| = 0$ in $C[0, |T|]$ by Stone-Weierstrass Theorem. It is clear $(V - U)P_n(|T|) \in \mathcal{K}(\mathcal{H})$ for all n . It follows that $(V - U)|T|^{1/2} \in \mathcal{K}(\mathcal{H})$ and therefore $|T|^{1/2}(V - U)|T|^{1/2} \in \mathcal{K}(\mathcal{H})$.

Then $T + K - \tilde{T} = |T + K|^{1/2}V|T + K|^{1/2} - |T|^{1/2}U|T|^{1/2}$

$$= |T + K|^{1/2}V(|T + K|^{1/2} - |T|^{1/2}) + (|T + K|^{1/2}(V - U)|T|^{1/2})$$

$$+ (|T + K|^{1/2} - |T|^{1/2})U|T|^{1/2} \in \mathcal{K}(\mathcal{H})$$

since $|T + K|^{1/2} - |T|^{1/2} \in \mathcal{K}(\mathcal{H})$.

Proposition 2.2

Let $T \in \mathcal{B}(\mathcal{H})$. Then $W_e(\tilde{T}) \subseteq W_e(T)$.

Proof

It is known that $W_e(T) = \bigcap \{\overline{W(T + K)} : K \in \mathcal{K}(\mathcal{H})\}$, that is, $W_e(T) = W_e(T + K)$ for all $K \in \mathcal{K}(\mathcal{H})$. By [7], we have $W_e(\tilde{T}) = W_e(T + K) \subseteq \overline{W(T + K)} \subseteq \overline{W(T + K)}$.

Thus $W_e(\tilde{T}) \subseteq \bigcap \{\overline{W(T + K)} : K \in \mathcal{K}(\mathcal{H})\} = W_e(T)$.

Lemma 2.3

Let $T \in \mathcal{B}(\mathcal{H})$. Then $\sigma_e(\tilde{T}) = \sigma_e(\tilde{T}^{(*)})$.

Proof

Recall that $\sigma_e(T) = \sigma_{le}(T) \cup \sigma_{re}(T)$. We first prove that $\sigma_{le}(\tilde{T}) \setminus \{0\} = \sigma_{le}(\tilde{T}^{(*)}) \setminus \{0\}$.

Suppose $\lambda \in \sigma_{le}(\tilde{T}) \setminus \{0\}$. Then there exists a unit vector sequence $\{x_n\}$ in \mathcal{H} such that $\{x_n\}$ converges weakly to zero and $\lim_{n \rightarrow \infty} \|(\tilde{T} - \lambda)x_n\| = 0$, that is, $\lim_{n \rightarrow \infty} \|(U^*\tilde{T}^{(*)}U - \lambda)x_n\| = 0$. In fact, we may choose $\{x_n\}$ in $N(T)^\perp$ such that $\|Ux_n\| = 1$ for all integer n . It follows that $\lim_{n \rightarrow \infty} \|(\tilde{T}^{(*)} - \lambda)Ux_n\| = \lim_{n \rightarrow \infty} \|(U^*\tilde{T}^{(*)} - \lambda)Ux_n\| \leq \lim_{n \rightarrow \infty} \|(\tilde{T} - \lambda)x_n\| = 0$.

Note that Ux_n converges weakly to zero, then $\lambda \in \sigma_{le}(\tilde{T}^{(*)})$. On the other hand, we can obtain $\sigma_{le}(\tilde{T}^{(*)}) \setminus \{0\} \subseteq \sigma_{le}(\tilde{T}) \setminus \{0\}$ by a similar method. Then $\sigma_{le}(\tilde{T}) \setminus \{0\} = \sigma_{le}(\tilde{T}^{(*)}) \setminus \{0\}$. It also follows that $\sigma_{re}(\tilde{T}) \setminus \{0\} = \sigma_{re}(\tilde{T}^{(*)}) \setminus \{0\}$ by the fact that $\sigma_{re}(A) = \sigma_{le}(A^*)$ for any $A \in \mathcal{B}(\mathcal{H})$. Then $\sigma_e(\tilde{T}) \setminus \{0\} = \sigma_e(\tilde{T}^{(*)}) \setminus \{0\}$. Next we show that $0 \in \sigma_e(\tilde{T})$ if and only if $0 \in \sigma_e(\tilde{T}^{(*)})$. This is equivalent to show that \tilde{T} is Fredholm if and only if $\tilde{T}^{(*)}$ is. Note that \tilde{T} (resp. $\tilde{T}^{(*)}$) is Fredholm if and only if U and $|T|^{1/2}$ (resp. $|T^*|^{1/2}$) are Fredholm. It follows that \tilde{T} is Fredholm if and only if $\tilde{T}^{(*)}$ is by the facts that $\tilde{T} = U^*\tilde{T}^{(*)}U$ and $\tilde{T}^{(*)} = U\tilde{T}U^*$. Above all, we have $\sigma_e(\tilde{T}) = \sigma_e(\tilde{T}^{(*)})$.

Theorem 2.4

Let $T \in \mathcal{B}(\mathcal{H})$. Then $W_e(\tilde{T}) = W_e(\tilde{T}^{(*)})$.

Proof

We recall that $\tilde{T} = \begin{pmatrix} 0 & 0 \\ 0 & X \end{pmatrix}$ and $\tilde{T}^{(*)} = \begin{pmatrix} 0 & 0 \\ 0 & Y \end{pmatrix}$, where X and Y are unitarily equivalent. Then we easily have $W_e(X) = W_e(Y)$

$$W_e(X) \subseteq W_e(\tilde{T}) \subseteq (W_e(X) \cup \{0\})^\wedge \text{ and } W_e(Y) \subseteq W_e(\tilde{T}^{(*)})$$

To complete the proof, it is sufficient to prove that $0 \in W_e(\tilde{T})$ if and only if $0 \in W_e(\tilde{T}^{(*)})$. Suppose $0 \in W_e(\tilde{T})$. If $0 \notin W_e(\tilde{T}^{(*)})$, then $W_e(\tilde{T}^{(*)}) = W_e(Y)$ and $0 \notin \sigma_e(\tilde{T}^{(*)})$. By Proposition 2.1, we also have $0 \notin \sigma_e(\tilde{T})$. It follows that $N(T)$ is finite-dimensional and X is Fredholm. Then $W_e(\tilde{T}) = W_e(X) = W_e(Y) = W_e(\tilde{T}^{(*)})$. This is a contradiction. Thus, $0 \in W_e(\tilde{T}^{(*)})$. Conversely, if $0 \in W_e(\tilde{T}^{(*)})$, we similarly have $0 \in W_e(\tilde{T})$.

Results and discussion

In this section, we give the main results of our study. We begin with the following Lemma.

Lemma 3.1

Let $T \in \mathcal{B}(\mathcal{H})$. Then

$$W_0(T^*) = (W_0(T))^* = \{\lambda : \bar{\lambda} \in W_0(T)\}.$$

Proof

We may assume that $\|T\| = \|T^*\| = 1$. Suppose $\lambda \in W_0(T)$. Then there exists a unit vector sequence $\{x_n\}$ in \mathcal{H} such that $\lim_{n \rightarrow \infty} \|Tx_n\| = 1$ and $\lim_{n \rightarrow \infty} \langle Tx_n, x_n \rangle = \lambda$. Now we

$$\lim_{n \rightarrow \infty} |\langle x_n, x_n \rangle - \langle Tx_n, Tx_n \rangle| = \lim_{n \rightarrow \infty} |(1 - T^*T)x_n, x_n| = 0,$$

which implies

$$\lim_{n \rightarrow \infty} \|(1 - T^*T)^{1/2}x_n\|^2 = 0.$$

$$\text{Then } \lim_{n \rightarrow \infty} \|(1 - T^*T)x_n\| = 0 \text{ and}$$

$$\lim_{n \rightarrow \infty} \|T^*Tx_n\| = 1.$$

In

$$\text{particular, } \lim_{n \rightarrow \infty} \|Tx_n\| = 1.$$

Hence,

$$\lim_{n \rightarrow \infty} |\langle T^*Tx_n, Tx_n \rangle - \langle x_n, Tx_n \rangle| = \lim_{n \rightarrow \infty} |\langle (T^*T - 1)x_n, Tx_n \rangle|$$

$$\leq \lim_{n \rightarrow \infty} \|(T^*T - 1)x_n\| \|Tx_n\| = 0.$$

We now have

$$\lim_{n \rightarrow \infty} \langle T^*Tx_n, Tx_n \rangle = \lim_{n \rightarrow \infty} \langle x_n, Tx_n \rangle = \bar{\lambda}.$$

Here put $y_n = Tx_n / \|Tx_n\|$. Then $\{y_n\}$ is a unit vector

sequence, $\lim_{n \rightarrow \infty} \|T^*y_n\| = 1$ and

$$\lim_{n \rightarrow \infty} \langle T^*y_n, y_n \rangle = \bar{\lambda},$$

which implies that $\bar{\lambda} \in W_0(T^*)$. Then $(W_0(T))^* \subseteq W_0(T^*)$. By symmetry, we have $W_0(T^*) = (W_0(T))^*$.

Lemma 3.2

Let $T \in \mathcal{B}(\mathcal{H})$. Then $\|\tilde{T} - \lambda\| = \|\tilde{T}^{(*)} - \lambda\|$ for all $\lambda \in \mathbb{C}$.

Proof

Let $\tilde{T} = \begin{pmatrix} 0 & 0 \\ 0 & X \end{pmatrix}$ and $\tilde{T}^{(*)} = \begin{pmatrix} 0 & 0 \\ 0 & Y \end{pmatrix}$ with respect to the space

decomposition $\mathcal{H} = N(T) \oplus N(T)^\perp$ and

$H = N(T^*) \oplus N(T^*)^\perp$ respectively,

where $Y = U_0 X U_0^*$, and U_0 is unitary.

Let $\lambda \in \mathbb{C}$, then we

$$\text{have } \tilde{T} - \lambda = \begin{pmatrix} -\lambda & 0 \\ 0 & X - \lambda \end{pmatrix}, \tilde{T}^{(*)} - \lambda = \begin{pmatrix} -\lambda & 0 \\ 0 & Y - \lambda \end{pmatrix},$$

$$\text{and } Y - \lambda = U_0(X - \lambda)U_0^*.$$

$$\text{Then } \|X - \lambda\| = \|Y - \lambda\|.$$

Case 1 $N(T) \neq \{0\}$ and $N(T^*) \neq \{0\}$.

Then

$$\|\tilde{T} - \lambda\| = \max\{|\lambda|, \|X - \lambda\|\} = \max\{|\lambda|, \|Y - \lambda\|\} = \|\tilde{T}^{(*)} - \lambda\|.$$

Case 2 $N(T) = \{0\}$ and $N(T^*) = \{0\}$. In this

case, $\tilde{T} - \lambda = X - \lambda$, $\tilde{T}^{(*)} - \lambda = Y - \lambda$. Clearly

$$\|\tilde{T} - \lambda\| = \|X - \lambda\| = \|Y - \lambda\| = \|\tilde{T}^{(*)} - \lambda\|.$$

Case 3 $N(T) = \{0\}$ and $N(T^*) \neq \{0\}$.

Then $\tilde{T} - \lambda = X - \lambda$. It follows

that $\|\tilde{T} - \lambda\| = \|X - \lambda\|$. Next we show

that $\|\tilde{T}^{(*)} - \lambda\| = \|Y - \lambda\|$. Otherwise,

if $\|Y - \lambda\| < |\lambda|$, then $-\lambda \in \rho(Y - \lambda)$, which

implies that Y is invertible. It follows that X is

also invertible since X and Y are unitarily

equivalent. Thus $\tilde{T} = X$ is invertible and so

are T and T^* . However, $N(T^*) \neq \{0\}$. This is a

contradiction.

$$\text{Hence } \|Y - \lambda\| \geq |\lambda| \text{ and } \|\tilde{T}^{(*)} - \lambda\| = \|Y - \lambda\|.$$

$$\text{Therefore } \|\tilde{T} - \lambda\| = \|\tilde{T}^{(*)} - \lambda\|.$$

Case 4 $N(T) \neq \{0\}$ and $N(T^*) = \{0\}$. The

proof is similar to Case 3.

Remark 3.3

Let U be a non-unitary isometry on \mathcal{H} . It is known that $U = U_0 \oplus U_1$ from the von Neumann-Wold Decomposition Theorem, where U_0 is unitary and U_1 is a unilateral shift.

Theorem 3.4

Let $T \in \mathcal{B}(\mathcal{H})$. Then $W_0(\tilde{T} - \lambda) = W_0(\tilde{T}^{(*)} - \lambda)$ for all $\lambda \in \mathbb{C}$.

Proof

Let $\lambda \in \mathbb{C}$, we

$$\text{have } \tilde{T} - \lambda = \begin{pmatrix} -\lambda & 0 \\ 0 & X - \lambda \end{pmatrix} \text{ and } \tilde{T}^{(*)} - \lambda = \begin{pmatrix} -\lambda & 0 \\ 0 & Y - \lambda \end{pmatrix}$$

with respect to the space

decomposition $\mathcal{H} = N(T) \oplus N(T)^\perp$ and

$\mathcal{H} = N(T^*) \oplus N(T^*)^\perp$ respectively, where $X - \lambda$ and $Y - \lambda$ are unitarily equivalent.

Case 1 $N(T) = \{0\}$ and $N(T^*) = \{0\}$. In this case $\tilde{T} - \lambda = X - \lambda$, $\tilde{T}^{(*)} - \lambda = Y - \lambda$.

The result follows.

Case 2 $N(T) \neq \{0\}$ and $N(T^*) \neq \{0\}$.

If $\|Y - \lambda\| = \|X - \lambda\| = |\lambda|$,

then

$$W_0(\tilde{T} - \lambda) = (\{\lambda\} \cup W_0(X - \lambda))^\wedge = (\{\lambda\} \cup W_0(Y - \lambda))^\wedge = W_0(\tilde{T}^{(*)} - \lambda)$$
 by Lemma 3.1.

If $\|X - \lambda\| = \|Y - \lambda\| > |\lambda|$,

then $W_0(\tilde{T} - \lambda) = W_0(X - \lambda) = W_0(Y - \lambda) = W_0(\tilde{T}^{(*)} - \lambda)$ by Lemma 3.2.

If $\|X - \lambda\| = \|Y - \lambda\| < |\lambda|$,

then $W_0(\tilde{T} - \lambda) = \{-\lambda\} = W_0(\tilde{T}^{(*)} - \lambda)$ by Lemma 3.1 again.

Case 3 $N(T) = \{0\}$ and $N(T^*) \neq \{0\}$.

Then $\tilde{T} - \lambda = X - \lambda$ and $W_0(\tilde{T} - \lambda) = W_0(X - \lambda)$. We next prove that $\|Y - \lambda\| > |\lambda|$. Note that $\tilde{\lambda}\tilde{T} = \lambda\tilde{T}$. Without loss of generality, we may assume that $\lambda = 1$.

We have

$$\|\tilde{T}^{(*)} - 1\| = \|\tilde{T} - 1\| = \|X - 1\| = \|Y - 1\| \geq 1$$
 by Lemma 3.1 and our assumption.

In fact, if $\|\tilde{T}^{(*)} - 1\| = \|\tilde{T} - 1\| < 1$, then we have that \tilde{T} is invertible. Then so is T . This contradicts with the assumption of this case. If $\|Y - 1\| = 1$, then $\|\tilde{T} - 1\| = 1$. Note that U is non-unitary isometry and $|T|^{1/2}$ is injective with dense range since $N(T) = 0$ and $N(T^*) \neq 0$. Then for any unit vector $x \in \mathcal{H}$, we have $|\langle \tilde{T}x, x \rangle - 1| \leq 1$, which implies

$$\left| \left\| |T|^{1/2}x \right\|^2 \left\langle U \frac{|T|^{1/2}x}{\| |T|^{1/2}x \|}, \frac{|T|^{1/2}x}{\| |T|^{1/2}x \|} \right\rangle - 1 \right| \leq 1.$$

$$\text{It is clear that } \left\langle U \frac{|T|^{1/2}x}{\| |T|^{1/2}x \|}, \frac{|T|^{1/2}x}{\| |T|^{1/2}x \|} \right\rangle \in W(U).$$

Note that $|T|^{1/2}$ has dense range. Then we can choose a unit vector $x_0 \in \mathcal{H}$ such

$$\left\langle U \frac{|T|^{1/2}x_0}{\| |T|^{1/2}x_0 \|}, \frac{|T|^{1/2}x_0}{\| |T|^{1/2}x_0 \|} \right\rangle \in (-1, 0)$$

that by Lemma 3.2. It follows that

$$\left| \left\| |T|^{1/2}x_0 \right\|^2 \left\langle U \frac{|T|^{1/2}x_0}{\| |T|^{1/2}x_0 \|}, \frac{|T|^{1/2}x_0}{\| |T|^{1/2}x_0 \|} \right\rangle - 1 \right| > 1.$$

This is a contradiction. Hence $\|Y - 1\| > 1$ and $W_0(\tilde{T}^{(*)} - 1) = W_0(Y - 1)$ by Lemma 3.2.

We then generally have $W_0(\tilde{T}^{(*)} - \lambda) = W_0(Y - \lambda) = W_0(X - \lambda) = W_0(\tilde{T} - \lambda)$ by Lemma 2.

Case 4 $N(T) \neq \{0\}$ and $N(T^*) = \{0\}$. The proof is similar to Case 3.

We recall that an inner derivation determined by $A \in \mathcal{B}(\mathcal{H})$ is defined by $\delta_A(X) = AX - XA$ for all $X \in \mathcal{B}(\mathcal{H})$. Stampfli in [8] gave the norm $\|\delta_A\|$ of δ_A by using of maximal numerical range, that is, $\|\delta_A\| = \inf \{\|T - \lambda\| : \lambda \in \mathbb{C}\}$. By Theorem 3, we have

Theorem 3.5

Let $T \in \mathcal{B}(\mathcal{H})$.

$$W_0(T) \subset \overline{W(\tilde{T})}.$$

If $\|T\| = \|\tilde{T}\|$, then $W_0(\tilde{T}) \subset W_0(T)$.

Proof

Without loss of generality, we may assume that $\|T\| = 1$.

Let $\lambda \in W_0(T)$, then there exists a unit vector sequence of $\{x_n\}$ in \mathcal{H} such that $\lim_{n \rightarrow \infty} \|Tx_n\| = 1$ and $\lim_{n \rightarrow \infty} \langle Tx_n, x_n \rangle = \lambda$, which implies

$$\lim_{n \rightarrow \infty} \| |T|^{1/2}x_n \| = 1 \text{ and } \lim_{n \rightarrow \infty} \|(1 - |T|)x_n\| = 0.$$

$$\lim_{n \rightarrow \infty} |\langle Tx_n, x_n \rangle - \langle \tilde{T}|T|^{1/2}x_n, |T|^{1/2}x_n \rangle|$$

$$= \lim_{n \rightarrow \infty} |\langle Tx_n, x_n \rangle - \langle U|T|x_n, |T|x_n \rangle|$$

$$= \lim_{n \rightarrow \infty} |\langle Tx_n, (1 - |T|)x_n \rangle|$$

$$\leq \lim_{n \rightarrow \infty} \|Tx_n\| \|(1 - |T|)x_n\| = 0.$$

Hence It follows that $\lim_{n \rightarrow \infty} \langle \tilde{T}|T|^{1/2}x_n, |T|^{1/2}x_n \rangle = \lambda$. Here put $y_n = |T|^{1/2}x_n / (\| |T|^{1/2}x_n \|)$. Then $\{y_n\}$ is a unit vector sequence and $\lim_{n \rightarrow \infty} \langle \tilde{T}y_n, y_n \rangle = \lambda$, and therefore $\lambda \in \overline{W(\tilde{T})}$.

We have $\|T\| = \|\tilde{T}\| = 1$.

Suppose $\lambda \in W_0(\tilde{T})$. Then there exists a unit vector sequence of $\{x_n\}$ in \mathcal{H} such that $\lim_{n \rightarrow \infty} \|\tilde{T}x_n\| = 1$ and $\lim_{n \rightarrow \infty} \langle \tilde{T}x_n, x_n \rangle = \lambda$.

It easily follows that $\lim_{n \rightarrow \infty} \| |T|^{1/2}x_n \| = \| |T|^{1/2} \| = 1$ and then $\lim_{n \rightarrow \infty} \|(1 - |T|)x_n\| = 0$. We easily have $\lim_{n \rightarrow \infty} \|(1 - |T|^3)x_n\| = 0$ also.

Thus $\lim_{n \rightarrow \infty} \|T|T|^{1/2}x_n\| =$
 $\lim_{n \rightarrow \infty} \langle |T|^3x_n, x_n \rangle = 1 = \|T\|.$

On the other hand,

$$\begin{aligned} & \lim_{n \rightarrow \infty} |\langle \tilde{T}x_n, x_n \rangle - \langle T|T|^{1/2}x_n, |T|^{1/2}x_n \rangle| \\ &= \lim_{n \rightarrow \infty} |\langle U|T|^{1/2}x_n, |T|^{1/2}x_n \rangle - \langle T|T|^{1/2}x_n, |T|^{1/2}x_n \rangle| \\ &= \lim_{n \rightarrow \infty} |\langle (U|T|^{1/2} - U|T|^{1/2})x_n, |T|^{1/2}x_n \rangle| \\ &= \lim_{n \rightarrow \infty} |\langle (U|T|^{1/2} - U|T|^{1/2})x_n, |T|^{1/2}x_n \rangle| \\ &\leq \lim_{n \rightarrow \infty} \|U|T|^{1/2}\| \|(1 - |T|)x_n\| \| |T|^{1/2}x_n \| = 0. \end{aligned}$$

Here put $y_n = |T|^{1/2}x_n / (\| |T|^{1/2}x_n \|).$

Then $\{y_n\}$ is a unit vector sequence and $\lim_{n \rightarrow \infty} \|Ty_n\| = \|T\| = 1$ and $\lim_{n \rightarrow \infty} \langle Ty_n, y_n \rangle = \lambda.$ Thus $\lambda \in W_0(T).$

Conclusions

If we let \mathcal{H} to be a complex separable Hilbert space and we let $\mathcal{B}(\mathcal{H})$ be the Banach algebra of all bounded linear operators on \mathcal{H} , we have characterized Aluthge transforms in Banach algebras. We have considered the classical and maximal numerical ranges of these transforms and finally we have given their relationships.

Conflicts of interest

Authors declare no conflict of interest.

References

- [1] Halmos PR. A Hilbert Space Problem Book, Springer Verlag New York; 1970.
- [2] Lumer G. Semi-inner product spaces. Trans Amer Math Soc 2016;100(5):29-43.
- [3] McIntosh A. Heinz inequalities and perturbation of spectral families, Macquarie Mathematics Reports 2006;291(2):79-86.

- [4] Jung IB, Ko E, Pearcy C. Aluthge transforms of operators. Integral Equations and Operator Theory 2000; 37(3):437-48.
- [5] Taylor J. A joint spectrum for several commuting operators. J Funct Anal 2019;6(2):172-91.
- [6] Okelo NB. On Characterization of Various Finite Subgroups of Abelian Groups. International Journal of Modern Computation, Information and Communication Technology 2018;1(5):93-8.
- [7] Okelo NB. On Normal Intersection Conjugacy Functions in Finite Groups. International Journal of Modern Computation, Information and Communication Technology 2018;1(6):111-5.
- [8] Okwany I, Odongo D, and Okelo NB. Characterizations of Finite Semigroups of Multiple Operators. Int J Mod Comput Info and Commun Technol Int J Mod Comput Info and Commun Technol 2018; 1(6):116-20.
- [9] Ramesh R, Mariappan R. Generalized open sets in Hereditary Generalized Topological Spaces J Math Comput Sci 2015;5(2):149-59.
- [10] Wanjara AO. On the Baire's Category Theorem as an Important Tool in General Topology and Functional Analysis. International Journal of Modern Computation, Information and Communication Technology 2019;2(4):27-31.
